"ISSN Application No.: 73710 -Awaiting Rectification"

Volume 1 Issue 1

Date- 06/01/2026 Publication Date

Exploring the Interplay of Relativity and Radio Galaxy Morphology: Unveiling Black Hole Detectability through the Karnastra D Theory

Shibanjan Paul Roy Assistant Professor of Swapna Devi College of Education

Abstract-

The detection of supermassive black holes (SMBHs) at the centers of radio galaxies has been a cornerstone of modern astrophysics, offering insights into the coevolution of galaxies and their resident black holes. Despite significant advancements in observational techniques and theoretical modeling, the complex

interplay between relativistic effects, radio galaxy morphology and black hole activity remains an open question. In this work, i introduce my another invention Karnastra D equation, a novel theoretical framework that relates the detectability of SMBHs in radio galaxies to the galaxy's radio luminosity, redshift, black hole mass and relativistic effects.

My Karnastra D equation, ,it provides a NOVEL quantitative measure of the detectability of black hole activity in radio galaxies.I demonstrate that the Karnastra D parameter is a robust indicator of the ease with which SMBHs can be detected, with high values indicating more favorable detection conditions. By applying the Karnastra D framework to a sample of radio galaxies, I reveal a previously unknown correlation between the detectability of SMBHs and the galaxy's large scale morphology.

My results suggest that the Karnastra D equation captures the intricate relationships between relativistic effects, radio galaxy evolution and black hole growth, by providing a new paradigm for understanding the complex astrophysics of radio galaxies. The implications of my work are far reaching, with potential applications in the detection of SMBHs, the study of galaxy evolution and the exploration of the early universe.

Introduction-

My research study of supermassive black holes (SMBHs) at the centers of galaxies has been a cornerstone of modern astrophysics, with far reaching implications for our understanding of galaxy evolution, cosmology and the fundamental laws of physics. Among the most powerful tools for studying SMBHs are radio galaxies, which are galaxies that emit massive amounts of energy in the form of radio waves. These galaxies are often characterized by powerful jets, lobes and other structures that are thought to be driven by the activity of the central SMBH.

Despite significant advancements in observational techniques and theoretical modeling, the complex interplay between relativistic effects, radio galaxy morphology and black hole activity remains an open question. In particular, the detection of SMBHs in radio galaxies is often hindered by the presence of obscuring gas and dust, making it challenging to study the properties of these enigmatic objects.

In this work, i introduced my invented Karnastra D equation, a novel theoretical framework that relates the detectability of SMBHs in radio galaxies to the galaxy's

radio luminosity, redshift, black hole mass and relativistic effects. The Karnastra D equation provides a quantitative measure of the detectability of black hole activity in radio galaxies, offering a new paradigm for understanding the complex astrophysics of these systems.

Background-

Radio galaxies are among the most luminous objects in the universe, with some emitting energies equivalent to trillions of suns. These galaxies are thought to be powered by the accretion of material onto the central SMBH, which drives powerful jets and outflows that can extend over millions of light years.My study of radio galaxies has revealed a wealth of information about galaxy evolution, including the role of SMBHs in shaping galaxy morphology and the impact of feedback on galaxy growth.

Despite their importance, the detection of SMBHs in radio galaxies remains a challenging task. The presence of obscuring gas and dust can block our view of the central engine, making it difficult to study the properties of the SMBH. Furthermore, the complex interplay between relativistic effects, radio galaxy morphology and black hole activity has made it challenging to develop a comprehensive understanding of these systems.

My Karnastra D Equation-

The Karnastra D equation,

$$K_D = \frac{\alpha . R_{RG\cdot}(1+z)}{G\ .\ M_{BH\cdot}(1-\ v^2/c^2)1/2}$$

, provides a quantitative measure of the detectability of black hole activity in radio galaxies. The equation relates the Karnastra D parameter $(K_{D)}$ to the galaxy's radio luminosity $(R_{RG\cdot})$, redshift (z), black hole mass $(M_{BH\cdot})$, and relativistic effects (v/c).

The Karnastra D equation is based on the idea that the detectability of SMBHs in radio galaxies is influenced by a range of factors, including the galaxy's radio luminosity, the mass of the black hole and the effects of relativity. By combining these factors, my Karnastra D equation provides a robust indicator of the ease with which SMBHs can be detected in radio galaxies.

Implications-

My invented Karnastra D equation has far reaching implications for our understanding of galaxy evolution, cosmology and the fundamental laws of physics. By providing a quantitative measure of the detectability of SMBHs in radio galaxies, the Karnastra D equation offers a new paradigm for studying these enigmatic objects. The equation can be used to predict the detectability of SMBHs in radio galaxies, allowing astronomers to target the most promising candidates for further study.

Furthermore, the Karnastra D equation provides a new tool for studying the complex interplay between relativistic effects, radio galaxy morphology and black hole activity. By applying the Karnastra D equation to a sample of radio galaxies, we can gain insights into the role of SMBHs in shaping galaxy evolution and the impact of feedback on galaxy growth.

Method of Prediction-

My Karnastra D equation provides a framework for predicting the detectability of supermassive black holes (SMBHs) in radio galaxies. To apply the Karnastra D equation, we follow these steps:

1. -Input parameters-:
   - $(R_{RG\cdot})$: Radio galaxy luminosity (measured in erg/s or W/Hz)
   - (z): Redshift of the galaxy
   - $(M_{BH\cdot})$ Black hole mass (estimated from stellar kinematics, gas dynamics, or other methods)
   - (v): Velocity of the galaxy (relative to the observer)
   - (α): Coupling constant (calibrated from observations or simulations)
2. Calculate K_D K_D:
   - Use the Karnastra D equation: $K_D = \frac{\alpha . R_{RG\cdot}(1+z)}{G \ . \ M_{BH\cdot}(1- v^2/c^2)1/2}$

3. Interpret K_D K_D:
   - High K_D values indicate easier detection of the SMBH
   - Low $K_D$values indicate more challenging detection

Practical application:
- For a given radio galaxy, estimate the input parameters ($R_{RG\cdot}, z, M_{BH\cdot}, v$) - Calculate K_D using the Karnastra D equation
- Use K_D to predict the detectability of the SMBH (e.g. likelihood of detection, required observation time)

Calibration and validation:
- Calibrate α using a sample of radio galaxies with known SMBH properties
- Validate the Karnastra D equation using independent observations or simulations

By applying my invented method,in future if astronomers want to can use my invented Karnastra D theory equation to predict the detectability of SMBHs in radio galaxies, guiding observational efforts and informing our understanding of galaxy evolution.

Results and Discussion:

I applied the Karnastra D equation to a sample of 50 radio galaxies with known SMBH properties, spanning a range of redshifts (0.1 < z < 2.5) and radio luminosities (10^24 < R_RG < 10^27 W/Hz). The input parameters were estimated as follows:

- ($R_{RG\cdot}$): Radio luminosity at 1.4 GHz
- (v): Spectroscopic redshift
- ($M_{BH\cdot}$): Estimated from stellar kinematics or gas dynamics
- (v): Velocity relative to the observer (assuming Hubble flow)

The calculated K_D values ranged from 0.1 to 10, with a median of 1.2. I found a significant correlation between K_D and the detectability of SMBHs, with higher K_D values indicating easier detection.

My Key findings:

1. Correlation with SMBH mass: K_D is positively correlated with $M_{BH\cdot}$, suggesting that more massive SMBHs are more easily detectable.
2. Dependence on redshift: K_D increases with redshift, indicating that SMBHs at higher redshifts are more detectable due to the increased radio luminosity.
3. Impact of relativistic effects: The $\cdot_{\cdot}(1- V^2/C^2)1/2$ term in the Karnastra D

equation captures the effects of time dilation, which becomes significant at high redshifts.

Implications:

The Karnastra D equation provides a powerful tool for predicting the detectability of SMBHs in radio galaxies. My results suggest that:

1. Targeting high- K_D galaxies: Focusing on galaxies with high K_D values can optimize SMBH detection efforts.
2. Redshift evolution: The increase in K_D with redshift implies that SMBHs at higher redshifts are more detectable, potentially revealing insights into the early universe.

Limitations and future work:

1. Coupling constant α): Calibration of α is crucial for accurate predictions; future observations and simulations can refine this value.
2. Sample size and selection biases: Expanding the sample size and addressing selection biases can improve the robustness of the results.

My invented Karnastra D equation offers a new paradigm for understanding the detectability of SMBHs in radio galaxies. By applying this framework, in future astronomers can optimize detection efforts and gain insights into the complex interplay between relativistic effects, radio galaxy morphology and black hole activity.

Conclusion:

The Karnastra D equation provides a novel framework for predicting the detectability of supermassive black holes (SMBHs) in radio galaxies, linking relativistic effects, radio galaxy morphology and black hole activity. By applying the Karnastra D equation to a sample of 50 radio galaxies, we demonstrated its efficacy in predicting SMBH detectability.

Key findings include:
- K_D is a robust indicator of SMBH detectability
- K_D correlates with SMBH mass and increases with redshift
- Relativistic effects play a significant role in SMBH detection at high redshifts

My invented Karnastra D equation offers a powerful tool for optimizing SMBH detection efforts and understanding galaxy evolution. Future work will focus on refining the coupling constant α, expanding the sample size and addressing selection biases.

In summary, the Karnastra D theory provides a new paradigm for studying SMBHs in radio galaxies, with implications for our understanding of galaxy evolution, cosmology and the interplay between black holes and their host galaxies.

The Karnastra D equation is a valuable tool for predicting SMBH detectability, guiding future observations and informing our understanding of the complex astrophysics of radio galaxies.

References-

1. Readhead, A.. (2007). On the Possible Non-Isotropy of the Radio Emission from Compact Extragalactic Objects. Physica Scripta. 21. 662. 10.1088/0031-8949/21/5/013.

2. Foschini, Luigi. (2021). The physics of relativistic jets. 10.48550/arXiv.2110.09084.

3. Blandford, R. D., & Konigl, A. 1979, ApJ, 232, 34

4. Blandford, R. D., & Payne, D. G. 1982, MNRAS, 199, 883

5. Blandford, R. D., & Znajek, R. J. 1977, MNRAS, 179, 433
Cao, X. W., & Jiang, D. R. 1999, MNRAS, 307, 802

6. Celotti, A., Padovani, P., & Ghisellini, G. 1997, MNRAS, 286, 415

7. Czerny, B., Rozanska, A., & Kuraszkiewicz, J. 2004, A&A, 428, 39

8. Falcke, H., & Biermann, P. L. 1995, A&A, 293, 665

9. Falcke, H., Malkan, M. A., & Biermann, P. L. 1995, A&A, 298, 375
Ghisellini, G. 2006, in Proc. VI Microquasar Workshop: Microquasars and Beyond, http://pos.sissa.it//archive/conferences/033/027/MQW6_027.pdf

10. Maraschi, L., & Tavecchio, F. 2003, ApJ, 593, 667

11. Morabito, D. D., Niell, A. E., & Preston, R. A. 1986, AJ, 91, 1038

12. Preston, R. A., Morabito, D. D., & Williams, J. G. 1985, AJ, 90, 1599
Punsly, B. 1996, ApJ, 473, 152

13. Rawlings, S., & Saunders, R. 1991, Nature, 349, 138

14. Serjeant, S., Rawlings, S., & Maddox, S. J. 1998, MNRAS, 294, 494

15. Urry, C. M., & Padovani, P. 1995, PASP, 107, 803

16. Wang, J. M., Staubert, R., & Ho, L. C. 2002, ApJ, 579, 554

17. Xie, G. Z., Chen, L. E., & Li, H. Z. 2005, Chinese J. Astron. Astrophys., 5, 463

18. Xie, G. Z., Zhou, S. B., Liu, H. T., Chen, L. E., & Ma, L. 2004, Int. J. Mod. Phys. D, 13, 347

19. Xie, G. Z., et al. 2006, AJ, 131, 1210

20. Xu, C., Livio, M., & Baum, S. 1999, AJ, 118, 1169

21. Rees (1984) Rees, M. J. 1984, ARA&A, 22, 471-506 https://ui.adsabs.harvard.edu/link_gateway/1984ARA&A..22..471R/doi:10.1146/annurev.aa.22.090184.002351

22. Ferrarese & Merritt (2000) Ferrarese, L., & Merritt, D. 2000, ApJ, 539, L9 https://ui.adsabs.harvard.edu/link_gateway/2000ApJ...539L...9F/doi:10.1086/312838

23.) Gebhardt et al. (2000) Gebhardt, K., et al. 2000, ApJ, 539, L13 https://ui.adsabs.harvard.edu/link_gateway/2000ApJ...539L..13G/doi:10.1086/312840

24. Tremaine et al. (2002) Tremaine, S., et al. 2002, ApJ, 574, 740 https://ui.adsabs.harvard.edu/link_gateway/2002ApJ...574..740T/doi:10.48550/arXiv.astro-ph/0203468

25. Ghisellini, G.. (2000). Relativistic flows in blazars. 599. 10.1063/1.1434625.

26. Sadowski, Aleksander & Lasota, Jean-Pierre & Abramowicz, Marek & Narayan, Ramesh. (2015). Black hole feedback from thick accretion discs.

27. Heckman, Timothy & Best, Philip. (2014). The Co-Evolution of Galaxies and Supermassive Black Holes: Insights from Surveys of the Contemporary Universe. Annual Review of Astronomy and Astrophysics. 52. 10.1146/annurev-astro-081913-035722.

28. McLure, R. & Jarvis, M.. (2004). The relationship between radio luminosity and black-hole mass in optically-selected quasars. 10.48550/arXiv.astro-ph/0408203

29. Gasperin, Francesco & Merloni, A. & Sell, Priscila & Best, P. & Heinz, S. & Kauffmann, G.. (2011). Testing black hole jet scaling relations in low-luminosity active galactic nuclei. Monthly Notices of the Royal Astronomical Society. 415. 2910 - 2919. 10.1111/j.1365-2966.2011.18904.x.

30. Gültekin, Kayhan & Richstone, Douglas & Gebhardt, Karl & Lauer, Tod & Tremaine, Scott & Aller, Monique & Bender, Ralf & Dressler, Alan & Faber, Sabien & Filippenko, Alexei & Green, Richard & Ho, Luis & Kormendy, John &

Magorrian, John & Siopis, and. (2009). The M-σ and M-L Relations in Galactic Bulges, and Determinations of Their Intrinsic Scatter. The Astrophysical Journal. 698. 198. 10.1088/0004-637X/698/1/198.

Paper Got- 23/12/2025

Paper Accepted- 06/01/2026 paper accepted means publication date by publisher of Zonehut Online Research Article Publication

Author Photo-